



Treatment Planning

28 slides, about 1-1.5 hour

JK Mitchell, DDS, MEd

Learning Objectives


Phase


3 Rehab

- Be able to recognize, explain, and determine the restorative implications of stressed pulp syndrome
- 2. Know the meaning of the following terms and be able to explain their clinical implications: Perio plastic surgery, functional crown lengthening, esthetic crown lengthening, biologic width, altered passive eruption, mucogingival defects
- 3. Be able to name and explain the options for single tooth replacement and multiple tooth replacement.
- 4. Explain the difference between a **bound vs unbound space** and the restorative implications.
- 5. Explain to a patient where implants are best used with an RPD

Data Collection \rightarrow Tx Plan

Here's where we are in the process now...






Phase 3 Goals

Phase 3 Rehab

"Begin with the end in mind"

- Up to now, we have been talking about treatment that you would be doing no matter what...Now we are getting to the point where the real options come in to play.
- Our goal is easy to state: **Restore form, function, and esthetics**. There are just so many ways to get there! And so many pitfalls in getting it right! And it is hard to explain at this point because *you have not been exposed to many of these treatments*.
- So just try to get an idea of some of the options. As you learn more about these in your advanced classes, you can see how your choices here affect your Phase 2 choices as well.



Phase 3 Treatment

ORTHODONTICS ENDODONTICS PERIODONTICS FIXED PROSTHODONTICS REMOVABLE PROSTHODONTICS

Phase

3 Rehab

Generally, treatments in Phase 3 must be carefully sequenced. It is crucial that you get this right or you will end up giving folks care for free when you make a mistake.

- 1. Orthodontics done in this phase is a stand-alone treatment. Otherwise, it would be in Phase 2.
- 2. Consider *any* potential Endodontic treatment before Prosthodontics. Teeth that are being considered as abutments for fixed or removable prosthetics should be tested for vitality.
- 3. **Periodontal adjunctive care can be quite complex.** More later, but there are times when you will need to start the crown preparation so you know exactly where the margins will end before you do the periodontal surgery.
- 4. Generally, Fixed Prosthetics comes before Removable Prosthetics.

Orthodontics





- There is so much that Orthodontics can do for your patients-both for esthetics and function- that does not involve preparing teeth!
- Any time there is an Orthodontic as well as a Prosthodontic option, you owe it to your patient to offer it and explain the advantages/ disadvantages.
- Never assume they would not be interested because of age, finances, or any other reason if they are otherwise a good candidate (Perio and Caries well controlled).

Endodontics



- Stressed pulp syndrome describes a tooth that has been repeatedly insulted with caries, restoration, or trauma.
- Even if still vital, preparing a stressed pulp may be the final straw. Note the size and shape of the pulp chamber compared to adjacent/contralateral tooth as one consideration.
- Look at it carefully and think: Would it be better to do endo before Pros is done? Or is the risk and quality outcome if done before or after?

Notice how much the size of the pulp on #13 is reduced compared to #12? Considering this history of deep decay, even if the pulp tests vital, at least *consider* an endodontic evaluation if you were going to, say, use it as a FPD or RPD abutment.



Abou-Ross M, J Pros Dent '82

Phase 3 Rehab

Phase 3 Periodontal Surgery

 Our Phase 2 Perio treatment was focused on treating disease, which included:

- 1. Initial Therapy (scaling and root planing to remove calculus)
- 2. Re-evaluation
- 3. If necessary, surgery for access to the calculus you couldn't reach or to reshape damaged bone or graft for repair.
- In Phase 3, Periodontal surgery falls in the category called Periodontal Plastic Surgery, which can include:
 - Reshape the periodontal architecture for esthetic reasons
 - Reshape supporting bone as an adjunct to Prosthodontic care
 - Reshaping or repairing mucogingival defects

Usually, your disease control goals are met in Phase 2 before you commit to this care in Phase 3, but to avoid duplicate surgery, you may choose to do both Phase 2 and Phase 3 surgery at the same time.

Periodontal Plastic Surgery

Phase 3 Rehab

Restorative Crown Lengthening



Before



After



There are two general types of crown lengthening procedures:

1. Restorative Crown Lengthening

- Generally, the body rejects restorative material placed too close to bone- you need about 2-3 mm between restoration and the crest of bone.
- This is called the Biologic Width. Putting a margin too close is a "biologic width violation"- it leads to chronic inflammation (redness, swelling, bleeds easily)
- These procedures reduce bone height so a restoration can be placed without violating the biologic width.

2. Esthetic Crown Lengthening

- Reshapes tissue for esthetics- to get a nice, even gingival architecture
- Can be done with Prosthodontic treatment or by itself.

Periodontal Plastic Surgery

- One common reason for Periodontal Plastic Surgery is "short teeth" or a "gummy smile" and one cause of that situation is Altered Passive Eruption. This happens when the teeth erupt, but the bone doesn't resorb correctly and the gingiva doesn't roll back, leaving the apparently "short teeth." Really, the teeth are the right size.
 - Teeth are amazingly consistent in size. Central incisors are nearly always at least 10 mm in height. If they are shorter than 10 mm, it's:
 - Worn down teeth- you can see the dentin
 - Excess gingiva -you'd have deep probing measurements
 - × Excess bone over the tooth.
 - You should be able to feel the CEJ with an explorer (gently!!! Get someone to show you how!). If not, then they haven't erupted correctly
- Doing prosthetics to "lengthen" the teeth is the WRONG treatment plan here!!! They would look like horse teeth, and the smile would still be "gummy"- the smile "frame" of lips is still in the same place.

For more on the "gummy smile" see J. William Robbins "Differential Diagnosis and Treatment of Excess Gingival Display, Pract Periodont Aesthet Dent 99: 11(2): 265-272



Phase

3 Rehab

This patient had an Esthetic Crown Lengthening procedure done to remove excess tissue left covering her normal sized teeth when altered passive eruption failed to expose them correctly.



Case 1: Altered Passive Eruption






Diagnosis

- 1. Short clinical teeth
 - < 10 mm central incisor
- 2. Can't feel CEJ with explorer clinically

Esthetic Crown Lengthening: Surgical Exposure



- Tissue reflected back to show location of bone.
- There needs to be 3 mm of space between the crest of bone and where you want the gingival crest to end up, and exactly the same shape.



Esthetic Crown Lengthening: Bone Recontouring



 Reshaping bone with chisels at the line angles.


Phase 3

Rehab

Esthetic Crown Lengthening: Bone Recontouring

Phase 3 Rehab

- Final position should be 2 mm apical to CEJ
- Notice how it's nicely shaped all the way to the line angles now



Esthetic Crown Lengthening: Tissue Placement

Phase 3 Rehab

- Tissue is replaced so that it is 3.0 mm from alveolar bone crest. This leaves 1 mm of tissue over enamel
- Without any further treatment, this is now a really nice esthetic result.
- Know when your patients might benefit from this so you can refer them to a Periodontist! (*Don't try this at home, folks...*)



Restorative Crown Lengthening

Phase 3 Rehab



Restorative Crown Lengthening



Before

After

I know what you're thinking- why not do the surgery first? Experience has taught us that only after you actually prepare the tooth do you know where the final margin needs to be. It's a real bummer to do the surgery, prep the tooth, and then have to re-do the surgery because you guessed wrong! In this case, the cusp has broken below the gingiva, so your margin needs to be that deep. But the bone is so close your margin would be impinging on the **biologic width**...what now?

- Go ahead and prepare your crown and place a provisional. That way, the surgeon knows exactly where the margin will be.
- 2. The surgeon then removes enough bone to leave the margin 3 mm from the height of bone.

Periodontal Plastic Surgery

Phase 3 Rehab

The frenum attachment has pulled the tissue away, leaving the exposed root and no attached gingiva in that area.

Case courtesy of Raleigh Periodontics and Implant Dentistry, Raleigh NC

Mucogingival defects can:

- Be unesthetic (on the maxillary)
- Cause sensitivity
- Predispose area to significant periodontal disease
- Periodontal grafts can restore the gingival tissue structure. This one was probably taken from the roof of the mouth and grafted into this area.
- Any time you are planning dentistry (margins, FPD clasps) in an area of a mucogingival defect, consider consulting with a periodontist.
- These defects trigger a D0180.

Case 2: Dennis



 Dennis is a 46 year old male who is concerned about the appearance of his teeth.

What is your first thought on the diagnosis?

- a Altered Passive Eruption
- b Worn Teeth

С

Congenitally Short Teeth

Rouen, France

A prosperous city under the Romans, Rotomagus was overun by the Vikings in the 800's. To stop the Viking raids, the French king gave them this part of France, which became known as Normandy ("north men land") with Rouen as the capital.


Although there has been a church on this site since the 300's, the current building was started in 1100 and finished in the 1300's after a fire.

During the Middle Ages, wool from England was woven into textiles and sold at the Champagne fairs, so it was very wealthy. A number of houses still exist from these days.

Break time!









Street market for local produce

The Gros Horologe (Big Clock) is from the late 1300's.






Street cafés are everywhere. Shops tempt you with pastry, or coffee, or chocolate shaped like a soccer ball...





The medieval center has been carefully preserved and rebuilt after WWII, including the Law Courts (above) and many of the wealthy merchant houses around the town square.





And Joan of Arc was burned in the square in Rouen during the Hundred Years War. The spot is marked and this statue is a few steps away.

Prosthodontics

FULL COVERAGE RESTORATIONS SINGLE TOOTH REPLACEMENTS MULTIPLE TOOTH REPLACEMENTS UNBOUNDED SPACE REPLACEMENTS

Full Coverage Restorations

- There are a number of materials to choose from....Consider each individual situation and use the best material for the situation
- You'll learn more about it in Fixed Pros and Esthetics this year and next.




Phase

3 Rehab

Single Tooth Replacement Options

Phase 3 Rehab



1. Nothing. Not all spaces have to be filled.

- Implant. Preferred replacement selection *but*:
 - □ Need enough bone (Width, height)
 - Need enough space (between teeth)
 - □ Pt health needs to be favorable

3. Fixed Partial Denture. Best choice:

- When implant not feasible
- When adjacent teeth need full coverage
- Removable Partial Denture. Usually a distant third in the replacement options. But, best choice:
 - When many other missing teeth in this arch
 - When significant bone and tissue missing as well. RPD can replace this with pink plastic for a better esthetic result.

Multiple Tooth Replacement Options

Phase 3 Rehab

- 1. Nothing. Again, not all spaces need to be restored.
- 2. Natural tooth FPD. Usually no more than two posterior or up to four anterior pontics per two abutment teeth.
- Implant FPD. Same issues as single tooth, but added complexity. Plan:
 - Max arch- One implant per tooth replaced
 - Mand arch- Two implants can support a three unit FPD
- **Removable Partial Denture**. If replacing more than two teeth, either implants or RPD's become the options, and since RPD's are, by a factor of 10, less expensive, they are a popular and very useful option.

Conventional FPDthis is a classic posterior 4 unit restoration



Implant FPD- planned for the mandible, we can fabricate a 3 unit FPD on 2 implants. On the maxilla, we would probably need 3 implants.

RPD- This appliance replaces the 4 incisors and 4 premolars by attaching to the remaining canines and molars.



Multiple Tooth Replacement Options: Unbounded Spaces

Phase 3 Rehab

Unbounded space

No tooth on the distal end of the space. Think it might be kind of loose and floppy?

Options?

Nothing, Implants, or RPD <u>+</u>implants



Functional tooth on either end of the space.


Options? Nothing, Implants, FPD, or RPD

 An "unbounded space" is one that does not have a tooth at one end.

- This, of course, rules out a conventional FPD- that used to mean an RPD was the only option.
- Now, implants offer a lot of other options, if the patient can afford them. But if not, RPD's can still work fine.
- BUT, generally, RPD's work better on the maxilla than the mandible. If the patient can only afford a few implants, put them in the mandible, either to eliminate or stabilize an RPD.

Multiple Tooth Replacement Options: Unbounded Spaces: Add Implants

Phase 3 Rehab



 An "unbounded space" is one that does not have a tooth at one end

This RPD was retrofitted with 2 implants at the distal end. Now it won't pull up with sticky food or wiggle loose with every tongue movement. a lot of


afford them. But if not, RPD's can still work fine.

BUT, RPD's work better on the maxilla than the mandible. If the patient can only afford a few implants, put them in the mandible, either to eliminate or stabilize an RPD.

Multiple Tooth Replacement Options: Unbounded Spaces- Add Implants

Phase 3 Rehab

This **maxillary RPD** is also filling unbounded spaces, but on the maxilla, with all the bone for stability and support, the implants were placed so that a metal clasp did not have to be placed in an area where it would show.



- **Combinations** of implants and removable prosthetic appliances can meet a variety of needs and financial means.
- **Gross oversimplification.** Prioritize implants as follows:
 - 1. Place implants to avoid having to place an RPD or FPD if possible
 - 2. Add implants to stabilize unbounded mandibular RPD
 - 3. Add implants to eliminate visible clasps, or stabilize maxillary RPD

The Edentulous Arch

And finally, there is the restoration of the **edentulous** arch.

- Nothing is still an option. Not a great one...but it will surprise you how well some people do.
- 2. Implant supported FPD. Great, but expensive
- 3. Implant supported RPD. Good option, less expensive.
- 4. Conventional complete denture. Least expensive.

There are some other twists on these choices, but these are the general concepts.



Maxillary Implant supported

RPD.

With at least 4 implants, the denture can be "palateless".

Notice that the esthetics of the

smile *may* be better with the RPD than the FPD because of the

shape of the teeth and the gingiva.

Maxillary Implant supported FPD



Phase

3 Rehab







Conventional complete dentures

Now that you've got the basics...

- Phase 3- Read textbook pages 200-211, not so much for the details, but for the overall concepts of when and why different choices are made.
- Maintenance- Read textbook pages 219-224. Be sure and read the "What's the Evidence" and "In Clinical Practice" boxes.
- Esthetics- Read articles on Esthetics in your D2L site, to a depth that you understand the basic steps of how to do an esthetic evaluation of a patient. Look at the clinic Esthetic Evaluation chart to get an appreciation (don't study it!) of what sort of things you will do in clinic.


In the Sophomore year, you are going to be learning how to make all these appliances- crowns, FPD's, RPD's, and complete dentures. There will be a lot of information on when and why as well as how, so knowing the framework of how this will fit into the patient's treatment will help you absorb these details so they will be usable in clinic, not just your next test.

Review

35

- 1 Recognize, explain, and determine the restorative implications of stressed pulp syndrome
 - Define the following terms and explain their clinical implications: Perio plastic surgery, Restorative crown lengthening, Esthetic crown lengthening, biologic width, altered passive eruption, mucogingival defects
- 3 List and explain, in terms a patient would understand, the options for single tooth replacement and multiple tooth replacement.
- 4 Explain the difference between a **bound vs unbound space** and explain how this affects your restorative options.
- 5 Explain to a patient where implants are best used with an RPD

2




Dennis: You chose a



Altered Passive Eruption

Altered passive eruption is diagnosed by clinically short teeth that are NOT worn on the incisals. These teeth are visibly worn, so this is probably not the best diagnosis.

Try again!



Dennis: You chose b


38



Worn Teeth

Correct! If you look at the incisal edges, you can see the facial and lingual edges of enamel with the dentin cupped out where it wears more quickly in between.

How do you fix it? *It's complicated....*



Dennis: You chose c

39



Congenitally short teeth

No, teeth are generally quite consistent. There *are* congenitally short teeth, but they are rare, and usually overall misshapen.

Try again!

